

## LABEL INFORMATION: ASPIRIN TABLETS



### Drug Facts

---

#### Active ingredient (in each tablet)

Aspirin 325 mg (NSAID\*) .....Pain reliever/fever reducer

\* nonsteroidal anti-inflammatory drug

#### Purposes

---

**Uses** ■ temporarily relieves pain due to: ■ headache ■ muscular aches ■ colds ■ minor pain of arthritis  
■ temporarily reduces fever

---

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

---

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

---

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

---

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for diabetes, gout, or arthritis

---

#### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

---

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

---

#### Directions

- adults: take 2 tablets every 4 hours, not more than 12 tablets in 24 hours
- children under 12 years: ask a doctor

---

#### Other information

- store at 15-25°C (59-77°F)
- avoid excessive heat above 40°C (104°F)

---

**Inactive ingredients** hydroxypropyl methylcellulose, polyethylene glycol, propylene glycol, starch

---

**Questions?** 1-800-841-8417 or zeemedical.com