

LABEL INFORMATION: PAINAID® BRF BACK RELIEF FORMULA TABLETS



Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 250 mg	Pain reliever
Magnesium salicylate tetrahydrate 250 mg (NSAID*)	Pain reliever

* nonsteroidal anti-inflammatory drug

Use temporarily relieves minor back pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Magnesium salicylate may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma, high blood pressure, heart disease, liver disease, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use salicylates during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- adults: take 2 tablets every 6 hours, not more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** magnesium 17 mg
- store at 15-25°C (59-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cornstarch, D&C yellow #10, FD&C yellow #6, hydroxypropyl methylcellulose, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide

Questions? 1-800-841-8417 or zeemedical.com