

LABEL INFORMATION: HISTENOL-FORTE® II MULTI-SYMPTOM COLD TABLETS



Drug Facts

Active ingredients (in each tablet)

Acetaminophen 325 mg	Pain reliever
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purposes

- Uses**
- temporarily relieves the following cold and flu symptoms:
 - nasal and sinus congestion
 - headache
 - minor aches and pains
 - helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- liver disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- adults: take 2 tablets every 4 hours, not more than 12 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 15-25°C (59-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C red #40, maltodextrin, microcrystalline cellulose, povidone, silica, sodium starch glycolate, starch, stearic acid

Questions? 1-800-841-8417 or zeemedical.com