



LABEL INFORMATION: IBUTAB™ ADVANCED PAIN RELIEF TABLETS

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID*)Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - headache ■ the common cold ■ muscular aches ■ backache ■ toothache ■ minor pain of arthritis
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ rash ■ skin reddening ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older ■ have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition
■ taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours
- children under 12 years: ask a doctor

Other information ■ store at 15-25°C (59-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients carnuba wax*, cellulose, croscarmellose sodium*, D&C yellow #10*, FD&C blue #2*, FD&C red #40*, FD&C yellow #6*, hydroxypropyl methylcellulose, lactose*, magnesium stearate, polydextrose, polyethylene glycol, polysorbate 80*, povidone*, red iron oxide*, silicon dioxide, sodium starch glycolate*, starch*, stearic acid*, titanium dioxide, triacetin*

*contains one or more of these ingredients

Questions? 1-800-841-8417 or zeemedical.com