

# LABEL INFORMATION: PAINAID® HEADACHE / PAIN RELIEF TABLETS



## Drug Facts

### Active ingredients (in each tablet)

	Purposes
Acetaminophen 110 mg .....	Pain reliever/fever reducer
Aspirin 162 mg (NSAID*).....	Pain reliever/fever reducer
Caffeine 32.4 mg .....	Pain reliever aid
Salicylamide 152 mg .....	Pain reliever aid

\* nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves pain due to: ■ headache ■ muscular aches ■ colds ■ flu  
■ temporarily reduces fever

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 8 tablets in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** ■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn  
■ you have asthma, high blood pressure, heart disease, liver disease, or kidney disease  
■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for diabetes, gout, or arthritis

### Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better  
■ ringing in the ears or loss of hearing occurs ■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

■ adults: take 2 tablets every 6 hours, not more than 8 tablets in 24 hours ■ children under 12 years: ask a doctor

**Other information** ■ store at 15-25°C (59-77°F) ■ avoid excessive heat above 40°C (104°F)

### Inactive ingredients

cellulose, croscarmellose sodium\*, FD&C yellow #6, magnesium stearate\*, povidone\*, starch, stearic acid\*  
\*contains one or more of these ingredients

**Questions?** 1-800-841-8417 or zeemedical.com